

A Distance-Limited Imaging Survey of Sub-Stellar Companions to Solar Neighborhood Stars

Joseph C. Carson

Max-Planck-Institut für Astronomie, Königstuhl 17, 69117 Heidelberg, Germany

Kyle D. Hiner¹, Gregorio G. Villar III, Michael G. Blaschak, Alexander L. Rudolph
Physics Department, California State Polytechnic University, Pomona, 3801 W. Temple
Ave., Pomona, CA 91768, U.S.A.

and

Karl R. Stapelfeldt

Jet Propulsion Laboratory, 4800 Oak Grove Dr., Pasadena, CA 91109, U.S. A.

Received _____; accepted _____

To be submitted to the Astronomical Journal.

¹present address: Department of Physics and Astronomy, University of California, Riverside, 900 University Ave., Riverside, CA 92507, U.S.A.

ABSTRACT

We report techniques and results of a Palomar 200-inch (5 m) adaptive optics imaging survey of sub-stellar companions to solar-type stars. The survey consists of K_s coronagraphic observations of 21 FGK dwarfs out to 20 pc (median distance ~ 17 pc). At $1''$ separation (17 projected AU) from a typical target system, the survey achieves median sensitivities 7 mag fainter than the parent star. In terms of companion mass, that corresponds to sensitivities of $50M_J$ (1 Gyr), $70M_J$ (solar age), and $75M_J$ (10 Gyr), using the evolutionary models of Baraffe and colleagues. Using common proper motion to distinguish companions from field stars, we find that no system shows positive evidence of a previously unknown substellar companion (searchable separation ~ 20 -250 projected AU at the median target distance).

Subject headings: methods: data analysis – stars: low-mass, brown dwarfs – surveys – techniques: high angular resolution

1. Introduction

The discovery of the brown dwarf GJ 229B (Nakajima et al. 1995) heralded a stream of direct detections of substellar objects. In particular, field surveys like the Sloan Digital Sky Survey (SDSS; Gunn & Weinberg 1995), the Two Micron All Sky Survey (2MASS; Skrutskie et al. 1997), and the Deep Near Infrared Survey (DENIS; Epchtein 1997) helped raise the number of brown dwarf (L and T type) identifications today to over 600 (Gelino et al. 2008). However the number of brown dwarfs identified as companions to main sequence stars remain few. At the time of this writing, there are only a handful of brown dwarfs confirmed as companions to main sequence stars. Brown dwarfs that are part of stellar systems are particularly interesting because they often yield insights into brown dwarf and planet formation around stars. For instance, statistics on the frequency of brown dwarf companions may shed light on the differences between planet, brown dwarf, and star formation. And unlike the case of discovered field brown dwarfs, the presence of a central star often reveals additional information on the presumably co-evolved brown dwarf – information like distance, metallicity, and age.

A number of high-contrast surveys have attempted to improve our knowledge of the moderate to wide separation (40AU to a few hundred AU) substellar companion population around stars. For example, Biller et al. (2007) and Metchev & Hillenbrand (2004) each surveyed samples (45 targets, 101 targets) of young (age $\lesssim 250$ Myr, $\lesssim 400$ Myrs), relatively nearby ($\lesssim 50$ pc, $\lesssim 160$ pc) stars using adaptive optics (AO) systems on the VLT and Palomar/Keck telescopes, respectively; Lowrance et al. (2005) used HST NICMOS to survey 45 young (median age ~ 150 Myr), nearby (average distance ~ 30 pc) stars for substellar companions; Lafrenière et al. (2007) used Gemini AO to observe 85 young (median age ~ 150 Myr), nearby (average distance ~ 22 pc) stars; Carson et al. (2005, 2006) used Palomar AO to survey 80 nearby (median distance ~ 17 pc) stars with unknown ages.

These references represent some of the larger direct-imaging high-contrast surveys, but there are a number of other surveys as well.

The observations described in this document largely provide an extension to Carson et al. (2005), although the new target list is focused more strongly on solar-type stars. (The Carson et al. [2005, 2006] survey looked mostly at K and M-dwarfs.) Most of the competing surveys (like Biller et al. 2007, Metchev & Hillenbrand 2004, Lowrance et al. 2005, Lafrenière et al. 2007, and others) have focused on observing the youngest nearby stars. While this allows for a maximal substellar-object self-luminosity (Baraffe et al. 2003), it inherently requires that one examines stars at somewhat larger distances, in order to achieve a large enough original sample to glean the youngest stars. Surveying only younger stars also leads to selection biases, as certain types of stars lend themselves better to age determinations than others (Mamajek & Hillenbrand 2008). For our survey, we avoid age requirements in an effort to achieve a more uniform census of the substellar companion population around the nearest solar-type stars.

While explorations of the substellar space around nearby FGK stars are scientifically interesting in their own right, they also provide important reconnaissance observations for the next generation of planet-search imaging surveys, like expected programs with Subaru (HiCIAO; Tamura et al. 2006), Palomar (Project 1640; Hinkley et al. 2008), VLT (SPHERE; Beuzit et al. 2008), Gemini (GPI; Macintosh et al. 2006), and potential space missions like Terrestrial Planet Finder Coronagraph (TPF-C)¹ and Terrestrial Planet Finder Interferometer (TPF-I)²/Darwin³. Information on the presence of brown dwarfs is important for these future planet surveys because the existence of an orbiting brown

¹<http://planetquest.jpl.nasa.gov/TPF-C>

²<http://planetquest.jpl.nasa.gov/TPF-I>

³<http://www.esa.int/science/darwin>

dwarf may affect the likelihood of there being a planet in that system. Even for southern hemisphere future planet searches, whose targets may not overlap with this document’s survey, the statistical information (from our survey and others) on brown dwarf frequencies may help guide southern surveys’ overall target selection strategy. In addition to information on orbiting brown dwarfs, future surveys will also benefit from our survey’s reports on discovered field objects close to nearby stars. Such information will ensure that future surveys do not spend unnecessary time following up these field objects for common proper motion tests. This information will also guide target selection, by providing information on field objects whose interfering light might adversely affect high-contrast sensitivities.

In the sections below we present techniques and results of our recently completed survey. Section 2 presents our target sample. Section 3 describes our observing techniques. In section 4 we present the data analysis techniques we used for this survey. In section 5 we summarize our survey sensitivities. Section 6 describes our results. We present our conclusions in section 7.

2. Targets

Our target selection process had its origins in work being carried out at Jet Propulsion Laboratory to select potential candidates for the proposed TPF-C mission (Levine et al. 2006). The main constraints for that selection process included FGK V spectral type, no known stellar multiplicity, closeness (<20 pc), and visible brightness ($V \lesssim 7$). For optimal observations from Palomar Observatory, we set a declination lower limit of -5 deg. We also removed any targets from our list that were already observed during the Cornell High-Order Adaptive Optics Survey (CHAOS; Carson et al. 2005) and other adaptive optics imaging programs, as determined by a standard literature search. We confirmed systems’ lack of multiplicity by on-telescope preliminary imaging. One target, GJ 564, had a previously

published brown dwarf binary companion (Potter et al. 2002). This fact eluded us in our initial literature search and preliminary imaging. Hence, we ended up observing this target and recording its current astrometry.

Our final target list consisted of 21 main sequence stars. This included 2 F stars, 14 G stars, and 5 K stars. All stars possessed well-characterized proper motion and parallax values as defined by *Hipparcos* (Perryman et al. 1997). As nearby stars, they typically possessed high proper motion (median target proper motion ~ 600 mas yr $^{-1}$), thus facilitating an efficient common proper motion follow-up strategy for candidate companions. A complete list of the target set is given in Table 1.

3. Observations

3.1. Coronagraphic Search Observations

To conduct our survey, we used the PALAO system (Troy et al. 2000) and the accompanying PHARO science camera (Hayward et al. 2001) installed on the Palomar 200 inch Hale Telescope. PALAO provided us with the high resolution (FWHM typically $\sim 0.14''$ in K_s and strehl ratio around 50%) necessary for resolving close companions. The accompanying PHARO science camera (wavelength sensitivity 1–2.5 μm and plate scale 40 mas pixel $^{-1}$) provided us with a coronagraphic imaging capability, along with a maximal field of view ($\sim 30''$). Our general observing strategy was to align the coronagraphic mask on a target star and take a series of K_s exposures, being careful to not saturate many pixels in the detector. (Occasionally, we saturated at the edges of the coronagraphic mask, where high noise levels already prevented any meaningful companion search.) We chose the K_s band because it provides high strehl ratios and a favorable relative flux between substellar companion and parent star. We planned our individual exposure times and number of

exposures per set to allow for no more than ~ 5 minutes (including overheads) between any target frame and the nearest sky image. This helped ensure that sky conditions did not significantly change between the target exposures and accompanying sky exposures. The sky exposures consisted of our taking, before and after the target sequence, and with the same setup as the target sequence, dithered images of a nearby empty sky region. We spent a comparable amount of time on sky as we did on target. We repeated the process of sky-target-sky as many times as was appropriate, with the goal of being able to detect (at 5-sigma) an ~ 18 -mag (K_s) object at $5''$ separation from the primary star. Once we completed the target/sky sets, we inserted a neutral density filter in the optical path and conducted dithered non-coronagraphic exposures of the target star. These images allowed us to characterize and record instrument and site observing conditions. Table 1 gives the relevant observing information for the individual targets.

3.2. Common Proper Motion Observations

For candidate companions detected in the previous procedures, we checked for physical companionship by using common proper motion observations. The nearby stars we observed tend to have high proper motions (on the order of a few hundred mas yr $^{-1}$). The vast majority of false candidate companions are background stars that tend to have very small proper motions compared to the parent star. Therefore, after recording our initial measurement, we waited for a timespan long enough for the parent star to move a detectable distance from the original position. In practice, this observable motion ended up being a minimum of ~ 3 pixels. After taking the second epoch observations, we checked whether the candidate maintained the same position with respect to the parent star. Target stars re-observed to check for common proper motion included GJ 895.4, 159, 230, 1095, 1085, 56.5, and 788. GJ 778 and 758 contained candidate companions, but were not re-observed

due to scheduling constraints (see discussion in Section 6).

4. Data Analysis

4.1. Reducing Images

We began our data reduction by median-combining each of the dithered sky sets. We then took each coronagraphed star image and subtracted the median-combined sky taken closest in time to the star image. (The typical separation in time between target and sky image was ~ 5.5 minutes.) We divided each of the sky-subtracted star images by a flat-field frame that we created, using standard procedures, from dark-frames and the dithered sky sets. We chose to use sky-flats, instead of the more conventional twilight or dome flats, in order to have a flat-field frame generated with an optical path most similar to that used with the target observation. We felt that this was important because we have seen transmission features (which we suspect reside on PHARO’s moveable optics) that change their observable position (by a few pixels perhaps) over the course of the night.

After performing the flat-fielding, we next median-combined each sequence of coronagraphic star frames. For this median combination, we used the images’ residual parent star flux (which leaked from around the coronagraph) to realign any frames that may have shifted because of instrument flexure. (Typical re-alignments were $\lesssim 1$ pixel.) We concluded that in-software re-alignment produced better overall sensitivities than just throwing-out mis-aligned frames. Next we applied a bad-pixel algorithm to remove suspicious pixels (defined as any pixel deviating from the surrounding 8 pixels by $\geq 5\sigma$) and replace them with the median of their 8 neighbors.

We next applied a Fourier filter to the resulting images to help remove non-point-like features such as unwanted internal instrument reflection and residual parent star flux. The

Fourier filter application entailed our multiplying a Fourier-transformed version of the final image by an exponential transmission function that minimized lower frequency signals. This Fourier filter, described in (Carson et al. 2005), has been shown to improve S/N by $\sim 25\%$ for a typical PHARO high-contrast image. Along with this S/N improvement, the typical point spread function (PSF) FWHM decreased by about 10% as a result of the Fourier filter application.

Finally, we investigated possibilities for taking advantage of the approximate symmetry of the coronagraphed PSF to self-subtract an inverted and/or rotated version of the PSF from the non-inverted image. We ended up deciding against using this technique as the final improvement was either marginal or non-existent.

4.2. Identifying Brown Dwarf Companions

Our first step in identifying candidate brown dwarf companions was to individually inspect each final Fourier-filtered and non-Fourier-filtered image for any potential companions. By choosing to examine both Fourier-filtered and non-Fourier-filtered final images, we effectively recognize that the filtering technique improves S/N in some instances and worsens it in others. For instance, Fourier-filtering works best in regions with shallowly sloping unwanted signal, like regions with internal instrument reflection. However, in other regions, the candidate companion S/N may suffer since the filtering always removes some true candidate companion flux.

The Palomar AO characteristic PSF “waffle pattern” (see Figure 1) is often seen as undesirable by observers, as the pattern degrades the potential PSF sharpness. While this is true in principle, the characteristic “waffle pattern”, in practice, provided an important way for us to distinguish true celestial objects (which had a well-patterned four-cornered

PSF) from statistical outliers in the image noise, which typically took on more arbitrary shapes. Thus, it provided an important first step in candidate companion identification.

While individual inspection of waffle patterns proved useful, we chose to also use an automated detection system to deliver more quantifiable sensitivity levels. Our automated algorithm operated by centering on every other pixel in the final reduced Fourier-filtered and non-Fourier-filtered image and creating there a $0.''16$ diameter flux aperture and $1.''2$ - $1.''6$ diameter sky annulus. After subtracting the background, the algorithm approximated a S/N level by dividing the measured aperture flux by the combined aperture flux Poisson noise and background noise; it approximated background noise from the standard deviation of the sky annulus pixels. In the end it outputted a final array with a S/N value for each sampled pixel. For each S/N map, there was also generated a map of measured background noise at each position (as estimated from the sky annuli). This outputted noise map essentially reflected the ability of the algorithm to detect (at a given thresh-hold S/N level) different brightness objects according to position on the array.

After generating maps for a given image, the program selected the S/N map pixel with the highest value, using a minimum value of five. It recorded the pixel position and then moved on to record the next highest S/N value greater than five. After each detection, it voided a $0.''4$ radius around the detected candidate object. This procedure continued until there were no more positions with S/N values greater or equal to five. (Of course, for many images, no positions possessed S/N levels greater than five.) After the algorithm identified the candidate sources, we re-examined the final images to ensure that the algorithm had indeed detected a true source as opposed to a systematic effect. Again, we searched for the Palomar adaptive optics signature “waffle pattern” to ensure a true physical source. We also made comparisons to images taken at other sources to ensure that the feature was indeed unique to the target image. In practice, we found that individual image inspection,

by eye, produced the most thorough identification of candidate companions. However, we felt that the automated detection was important as well, in order to provide quantifiable detection sensitivities, as well as a second-check for our visual inspections.

We acknowledge that the use of our automated detection routine has some drawbacks. Notably, there are instances where the algorithm over-estimates noise levels. For instance, close to the parent star PSF, the algorithm can mistake what may be a well-ordered parent star PSF slope for a random fluctuation in background noise. Additionally, the algorithm may also over-estimate the noise close to field stars; if a field star happens to fall in the sky annulus, the algorithm will determine that region to have excessively high background noise. Thus, only the brightest candidate objects would be detected near these field star positions. While these instances are not ideal, we conclude that is an acceptable compliment to our careful visual inspections. In Section 5 we discuss how we may generate limiting magnitudes and brown dwarf mass limits from these algorithm-generated noise maps.

In cases where we positively identified a potential brown dwarf companion, we next estimated its apparent K_s magnitude by comparing its flux to the non-coronagraphic parent-star calibration images and published 2MASS K -magnitudes (Skrutskie et al. 1997). Resulting magnitudes are displayed in Table 2. Once we established an apparent K_s magnitude, we derived a corresponding absolute K_s magnitude, assuming the candidate had a distance equal to the parent system. Thanks to observational surveys such as *Hipparcos* (Perryman et al. 1997), all of our parent stars had well-defined parallaxes and therefore distances. With an approximate absolute K_s magnitude in hand, we combined published brown dwarf observational data (Leggett et al. 2000, Leggett et al. 2002, Burgasser et al. 1999, Burgasser et al. 2000, Burgasser et al. 2002, Burgasser et al. 2003, Geballe et al. 2002, Zapatero et al. 2002, Cuby et al. 1999, Tsvetanov et al. 2000, Strauss et al. 1999, and Nakajima et al. 1995) with theoretical data from Baraffe et al. (2003) to extrapolate

constraints on the object’s mass. An object whose potential mass fell within acceptable brown dwarf restrictions was designated for common proper motion follow-up observations.

For our follow-up observations, we used Hipparcos published common proper motion values (Hipparcos catalogue; Perryman et al. 1997) to determine the expected movement of the parent system. Since background and field stars are unlikely to possess proper motions identical to the parent system’s, we used common proper motion as a strong support for a physical companionship. For candidate companions, we used the PSF central peak to identify position. For the obscured parent star, we used the waffle-pattern four corners (which resided outside the opaque coronagraph spot) to create a well-defined cross-hairs that revealed the central position. We could typically constrain the relative offset between parent star and candidate companion by fractions of a pixel, depending on S/N levels. Measuring the candidate companion’s relative position over the two epochs, we were able to distinguish physical companionships from chance alignments. We record positions in Table 2.

5. Survey Sensitivities

5.1. Determining Limiting Magnitudes

To quantify detection sensitivities from the algorithm-generated noise maps described in Section 4.2, we looked to determine the faintest detectable magnitude as a function of angular separation from each parent star. We began by sampling each noise map (including those deriving from Fourier-filtered and non-Fourier-filtered images) and selecting, for each pixel, the smaller of the two noise values. The resulting composite noise map array therefore reflected the best sensitivities from each of the two final images. Figure 2 displays a sample image sequence, where a Fourier-filtered and non-Fourier-filtered image are combined to

create a composite noise map.

For the composite noise map, we next determined the median values in a series of concentric $0''.20$ -thick rings centered on the noise map center. The median values therefore represented typical noise as a function of distance from the central star. For each noise value, we then determined the minimum apparent K_s -magnitude where signal exceeded the combined Poisson noise and ring noise by a factor greater or equal to 5; we were able to convert noise values (in units of detector counts) to K_s -magnitudes using parent star calibration data described in Section 3.1. In Figure 3 we plot resulting measurements for median survey sensitivities (middle curve), the best 10% of observations (lower curve), and the worst 10% of observations (top curve). Refer to Table 3 for a summary of minimum detectable magnitudes for each of the individual targets.

Another commonly used statistic for describing sensitivities for high-contrast companion surveys is the limiting differential magnitude as a function of angular separation from the parent star. In other words, how many times dimmer may a companion object be before we lose it in the parent star noise? Figure 4 plots differential magnitudes for median survey sensitivities as well as the best and worst 10% of observations.

5.2. Mass Sensitivities

Determining sensitivities according to companion mass is complicated by the fact that brown dwarfs of a given mass dim over time. Nonetheless, to get a general idea of detectable masses, we may assume different test ages and then use models by Baraffe et al. (2003) to transform our minimum detectable brightnesses into brown dwarf masses. Figure 5 shows a comparison of median sensitivities assuming 1 Gyr, solar age, and 10 Gyr target ages.

6. Results

After conducting all of our data analysis, we concluded that zero systems showed positive evidence of a brown dwarf companion (that was not previously known). We did re-detect the brown dwarf binary orbiting GJ 564, discovered by Potter et al. (2002). In total, we detected 48 field objects (including the binary brown dwarf) around 10 targets stars. The GJ 778 and GJ 758 fields both contained candidate companions, but were not re-observed for common proper motion follow-up tests due to scheduling constraints. 2MASS data (Skrutskie et al. 1997) reports K-band field star densities of ~ 12 stars and ~ 6 stars per PHARO field of view, for the respective GJ 778 and GJ 758 star neighborhoods. Given the relatively high field star densities for these regions, the chances of these candidates being field stars are high. In the end, however, we cannot confirm or reject that one of these candidates may be a brown dwarf companion. Instead, we simply report the photometry and astrometry for the detected objects. Table 2 presents the measurements for these objects as well as all the other field objects detected in our survey.

7. Discussion

As mentioned in the previous paragraph, our survey found no evidence of new brown dwarf companions, for orbital separations akin to our own outer solar system. However, even for targets with no candidate companions, we cannot rule out the possibility that one or more new brown dwarfs exist around the targeted stars, even at the semi-major axes for which our survey is most sensitive. For instance, a substellar companion near conjunction, in an orbit close to edge-on, may be impossible to resolve from the parent star PSF, regardless of the companion’s luminosity or semi-major axis. Furthermore, even for a bright brown dwarf with a face-on orbital inclination, a brown dwarf’s orbital eccentricity might lead to a range of possible projected separations, which could lead to a null detection.

Extracting rigorous companion statistics is therefore complicated by factors such as unknown orbital characteristics. For example, if typical brown dwarf orbits are highly eccentric, the typical semi-major axis regime that our survey covers is most likely narrower than the *projected* orbital separation we probe, as shown in orbital simulations presented in Carson et al. (2006). Furthermore, extracting the companion fraction for a given substellar mass range is complicated by the fact that one must assume a system age in order to translate the survey sensitivity floor (in terms of K_s mag) into a minimum detectable mass (see discussion in Section 5.2). Since most of our stars have unknown ages, to extract a companion fraction, one must resort to a statistical inference of target star ages, using a method such as galactic birth models (like that used in Burgasser 2004) or stellar metallicity relations (like that used in Carson et al. 2006). (Alternative age determination methods, such as those using Ca II emission, lithium abundance, and X-ray activity, provide poor constraints for target sets older than a couple hundred Myrs, and are therefore not useful for our applications.)

The extractable companion frequency also depends on the relative mass function of substellar companions. For example, even if we limit ourselves to a constrained mass range (like $20 M_J$ to $40 M_J$), the companion fraction uncertainties may depend sensitively on whether the majority of brown dwarfs resides near the lower boundary or the higher boundary of our mass range; if the majority resides near the lower mass boundary, there is a greater chance that our null result is due to limiting sensitivities, as opposed to a true lack of companions.

It is possible for one to make educated assumptions for all of the aforementioned factors, and then run detailed Monte Carlo population simulations to conclude a companion fraction (e.g. Nielsen et al. 2008, Carson et al. 2006). Published Monte Carlo population analyses have shown that a large target data set ($\gtrsim 60$ stars) is typically required to provide

meaningful results. For instance, Carson et al. (2006), using an 80 star sample, concludes a brown dwarf companion (25-100 AU semi-major axis) fraction of 0% to 9%. Nielsen et al. (2008) concludes, from a 60 star sample, a planet/brown-dwarf ($> 4 M_J$) companion (20-100 AU semi-major axis) fraction of 0% to 20%. Considering these relatively large uncertainties, and that our 21 star sample is significantly smaller than these other surveys, we believe that we cannot conclude meaningful companion fractions from our data set alone. We postpone, for a future paper, a detailed Monte Carlo population analysis that combines several surveys' data sets to derive the most meaningful companion fractions.

In addition to explorations of brown dwarf companions, our survey also reported astrometry and photometry for all detected field objects. The reporting of such objects may have potential benefits to future surveys, by providing possible reference star candidates, preventing future brown dwarf candidate false identifications, and yielding data on objects whose interfering light might impede future planet-search observations.

We thank the Palomar Observatory staff for their support of these observations. We thank our anonymous referee for useful comments. Part of the research conducted by J.C.C was supported by an appointment to the NASA Postdoctoral Program at the Jet Propulsion Laboratory, administered by Oak Ridge Associated Universities through a contract with NASA.

Facilities: Hale

REFERENCES

- Baraffe, I., Chabrier, G., Barman, T. S., Allard, F., & Hauschildt, P. H. 2003, *A&A*, 402, 701
- Beuzit, J.-L. et al. 2008, *SPIE*, in press
- Biller, B. A. et al. 2007, *ApJS*, 173, 143B
- Burgasser, A. J. 2004, *ApJS*, 155, 191B
- Burgasser, A. J., et al. 1999, *ApJ*, 522, 65
- Burgasser, A. J., et al. 2000, *AJ*, 120, 1100
- Burgasser, A. J. et al., 2002, *ApJ*, 564, 421
- Burgasser, A. J., McElwain, M. W., & Kirkpatrick, D. J., 2003, *AJ*, 126, 24 87
- Carson, J. C., Eikenberry, S. S., Brandl, B. R., Wilson, J. C., and Hayward, T. L. 2005, *AJ*, 130, 1212
- Carson, J. C., Eikenberry, S. S., Smith, J. J., and Cordes, J. M. 2006, *AJ*, 132, 1146C
- Cuby, J. G., Saracco, P., Moorwood, A. F. M., D’Odorico, S., Lidman, C., Comerón, F., & Spyromilio, J. 1999, *A&A*, 349, L41
- Geballe et al. 2002, *ApJ*, 564, 466
- Gelino, C. R., Kirkpatrick, J. D., & Burgasser, A. J. 2008, *DwarfArchives.org*, <http://dwarfarchives.org>
- Gliese, W., & Jahreiss, H. 1991, *The Third Catalogue of Nearby Stars: Errors and Uncertainties* (Heidelberg: Astron. Rechen-Institut)

- Gunn, J. & Weinberg, D. 1995, WFSD Conf, 3
- Hayward, T. L., Brandl, B., Pirger, B., Blacken, C., Gull, G. E., Schoenwald, J., & Houck, J. 2001, PASP, 113, 105
- Hinkley, S., Oppenheimer, B. R., Brenner, D., Parry, I. R., Sivaramakrishnan, A., Soummer, R., & King, D. 2008, SPIE, in press
- Lafrenière, D. et al. 2007, ApJ, 670, 1367L
- Leggett, S. K. et al. 2000, ApJ, 536, 35
- Leggett, S. K. et al. 2002, ApJ, 564, L452
- Levine, M., Shaklan, S., & Kasting, J. eds. 2006, Terrestrial Planet Finder Coronagraph Science and Technology Definition Team (STDT) Report, JPL document D-34923 (Pasadena: JPL)
- Lowrance, P. J. et al. 2005, AJ, 130, 1845
- Macintosh, B. et al. 2006, SPIE, 6272E, 18M
- Mamajek, E. E. & Hillenbrand, L. A. 2008, ApJ, in press
- Metchev & Hillenbrand 2004, ApJ, 617, 1330
- Nakajima, T., Oppenheimer, B. R., Kulkarni, S. R., Golimowski, D. A., Matthews, K., & Durrance, S. T., 1995, Nature, 378, 463
- Nielsen, E. L., Close, L. M., Biller, B. A., Masciadri, E., & Lenzen, R. 2008, ApJ, 674, 466
- Perryman, M. A. C. et al. 1997, A&A, 323, 49
- Potter, D., Martín, E. L., Cushing, M. C., Baudoz, P., Brandner, W., Guyon, O., & Neuhauser, R. 2002, ApJ, 567L, 133P

- Skrutskie, M. F., et al. 1997, in *The Impact of Large-Scale Near-IR Sky Surveys*, ed. F. Garzon (Dordrecht: Kluwer), 25
- Strauss, M. A. et al. 1999, *ApJ*, 522, 61
- Tamura, Motohide et al. 2006, *SPIE*, 6269E, 28T
- Troy et al. 2000, *SPIE*, 4007, 31
- Tsvetanov, Z. I. et al. 2000, *ApJ*, 531, 61
- Zapatero Osorio, M. R., Béjar, J. S., Martín, E. L., Rebolo, R., Barrado y Navascués, D., Mundt, R., Eislöffel, J. & Caballero, J. A. 2002, *ApJ*, 578, 536




Fig. 1.— Two example reduced K_s images of GJ 230, taken in November, 2007 using the Palomar 200-inch Adaptive Optics and accompanying PHARO science camera. The image on the left is a non-coronagraphic GJ 230 image taken with a neutral density filter. The slight elongation of the central PSF feature toward the lower left direction is an artifact of the neutral density filter, and does not occur in non-neutral-density-filter images. The image on the right was taken with no neutral density filter, but with a $0.''91$ opaque spot positioned over the star. The images illustrate PALAO’s characteristic AO-reconstructed PSF as seen in both coronagraphic and non-coronagraphic imaging.




Fig. 2.— GJ 230 Final Images and Noise Map. a) Left: GJ 230 coronagraphic final image. b) Center: GJ 230 coronagraphic, Fourier-filtered, final image. c) Right: a composite noise map created from (a) and (b). The greyish, slightly offset ring segments around the star in (a) and (b) result from internal instrument reflection.




Fig. 3.— K_s -band sensitivity curves displaying limiting magnitude as a function of separation from the parent star. The top curve represents the median sensitivities for the worst 10% of observations. The middle curve represents median survey sensitivities. The bottom curve represents median sensitivities for the best 10% of observations. “Best 10%” and “Worst 10%” are defined by a combination of parent star brightness, seeing conditions, and adaptive optics performance. All minimum magnitudes correspond to 5-sigma detections.




Fig. 4.— K_s -band sensitivity curves displaying limiting differential magnitude (K_s -companion minus K_s -parent) as a function of separation from the parent star. The middle curve represents median survey sensitivities. The top curve represents median sensitivities for the worst 10% of our data. The bottom curve shows the best 10%. Limits represent 5-sigma detections.




Fig. 5.— Minimum detectable mass as a function of projected separation for median target sensitivities and distances. We compare results for assumed 1 Gyr, solar age, and 10 Gyr targets, using evolutionary models by Baraffe et al. (2003). We derived these curves by first plotting minimum detectable K-short magnitude versus arcsecond separation for all targets. Next we median-combined the K-short curves to derive typical sensitivities (see Figure 3, middle curve). We transformed these K-short magnitudes into masses using Hipparcos distances and Baraffe et al. (2003) evolutionary models. Finally, we transformed our arcsecond axis to a projected AU separation using the median target distance.

Table 1. Nearby Star Target List

Parallax (mas)	Proper Motion		V (mag)	Name	Dates of Coronagraphic Observations	Net Exposure Time (sec)
	RA (mas yr ⁻¹)	Dec (mas yr ⁻¹)				
59.31	1106.1	113.0	6.76	GJ 895.4	2005 Dec; 2006 May; 2007 Nov	4916
52.00	151.2	-252.0	5.38	GJ 159	2006 Dec; 2007 Nov	581
55.20	78.1	-297.1	6.43	GJ 230	2005 Dec; 2007 Nov	2149
59.31	29.4	-186.1	5.54	GJ 1095	2006 Dec; 2007 Nov	586
59.46	-34.1	-34.45	7.17	GJ 56.5	2006 Dec; 2007 Nov	581
64.25	62.4	-230.7	5.97	GJ 1085	2006 Dec; 2007 Nov	595
59.52	-171.2	-1164.2	6.97	GJ 295	2006 Dec	340
57.57	-359.8	139.3	5.95	GJ 484	2006 Dec	283
56.92	468.5	296.8	5.91	GJ 788	2006 May; 2006 Dec	3054
64.71	-122.3	-103.3	6.76	GJ 227	2005 Dec; 2006 Dec	1982
52.25	-191.1	-115.4	5.95	GJ 334.2	2006 Dec	297
71.04	-315.9	55.2	5.03	GJ 407	2005 Dec; 2006 May	2046
56.82	223.7	-477.5	6.25	GJ 547	2006 May	1113
55.11	132.5	-298.4	6.61	GJ 614	2006 May	1338
64.54	82.0	162.9	6.37	GJ 758	2006 May	361
67.14	-529.2	-428.9	5.37	GJ 376	2006 May	2028
55.73	144.7	32.4	5.86	GJ 564	2006 May	487
69.61	-571.2	52.6	6.66	GJ 611	2006 May	476
55.37	123.5	854.7	6.76	GJ 651	2006 May	1427
64.17	-1002.8	-912.6	7.28	GJ 778	2006 May	1517
70.07	-27.7	87.9	5.63	GJ 311	2006 Dec	297

Note. — Parallax, proper motion, and V-magnitude are from Hipparcos (Perryman et al. 1997). All names follow the Gliese catalog system (Gliese & Jahreiss 1991).

Table 2. Detected Field Objects

Parent Field	ρ (arcsec) ^a	θ (deg) ^b	Observation	
			Date	K_s (mag)
GJ 895.4	9.619 ± 0.006	120.7 ± 0.3	2007 Nov	15.53 ± 0.16
GJ 895.4	5.375 ± 0.008	287.1 ± 0.3	2007 Nov	16.94 ± 0.17
GJ 895.4	12.992 ± 0.005	308.7 ± 0.3	2007 Nov	17.11 ± 0.12
GJ 895.4	12.724 ± 0.004	328.9 ± 0.3	2007 Nov	17.67 ± 0.11
GJ 895.4	11.629 ± 0.008	342.30 ± 0.3	2007 Nov	17.13 ± 0.07
GJ 895.4	11.544 ± 0.007	341.7 ± 0.3	2007 Nov	17.35 ± 0.09
GJ 895.4	9.771 ± 0.006	347.9 ± 0.3	2007 Nov	18.24 ± 0.12
GJ 895.4	6.572 ± 0.100	338.4 ± 0.3	2007 Nov	19.13 ± 0.13
GJ 895.4	16.279 ± 0.009	315.1 ± 0.3	2007 Nov	18.24 ± 0.11
GJ 895.4	3.370 ± 0.004	118.0 ± 0.4	2007 Nov	16.81 ± 0.08
GJ 159	15.747 ± 0.013	310.8 ± 0.3	2006 Dec	16.89 ± 0.02
GJ 230	16.508 ± 0.007	321.8 ± 0.3	2007 Nov	18.58 ± 0.04
GJ 1085	10.600 ± 0.007	148.1 ± 0.3	2006 Dec	17.12 ± 0.08
GJ 1085	13.892 ± 0.025	226.2 ± 0.3	2006 Dec	17.63 ± 0.12
GJ 1095	10.459 ± 0.006	84.4 ± 0.3	2006 Dec	15.79 ± 0.03
GJ 1095	9.491 ± 0.003	87.5 ± 0.3	2006 Dec	18.91 ± 0.34
GJ 56.5	9.510 ± 0.023	130.0 ± 0.3	2006 Dec	16.97 ± 0.06
GJ 788	7.263 ± 0.008	119.0 ± 0.3	2006 May	17.78 ± 0.20
GJ 788	7.096 ± 0.009	119.0 ± 0.3	2006 May	18.09 ± 0.2
GJ 564*	2.600 ± 0.019	101.3 ± 0.4	2006 May	–
GJ 564*	2.600 ± 0.012	102.6 ± 0.4	2006 May	–
GJ 758	11.078 ± 0.009	57.0 ± 0.3	2006 May	13.12 ± 0.03
GJ 758	7.216 ± 0.006	358.8 ± 0.3	2006 May	13.38 ± 0.04
GJ 758	10.396 ± 0.003	193.4 ± 0.3	2006 May	8.43 ± 0.04
GJ 758	10.279 ± 0.003	210.9 ± 0.3	2006 May	12.88 ± 0.03
GJ 758	14.297 ± 0.015	135.2 ± 0.3	2006 May	14.12 ± 0.03
GJ 758	11.131 ± 0.012	319.2 ± 0.3	2006 May	15.09 ± 0.10
GJ 758	15.862 ± 0.010	235.0 ± 0.3	2006 May	13.37 ± 0.04
GJ 778	9.778 ± 0.007	101.8 ± 0.3	2006 May	17.61 ± 0.02
GJ 778	8.143 ± 0.005	87.1 ± 0.3	2006 May	16.00 ± 0.02
GJ 778	6.421 ± 0.007	81.8 ± 0.3	2006 May	16.63 ± 0.04

Table 2—Continued

Parent Field	ρ (arcsec) ^a	θ (deg) ^b	Observation	
			Date	K_s (mag)
GJ 778	8.222 ± 0.008	310.3 ± 0.3	2006 May	14.81 ± 0.02
GJ 778	8.857 ± 0.010	301.3 ± 0.3	2006 May	13.52 ± 0.03
GJ 778	11.111 ± 0.006	263.7 ± 0.3	2006 May	13.61 ± 0.03
GJ 778	6.151 ± 0.008	227.3 ± 0.3	2006 May	15.21 ± 0.03
GJ 778	14.726 ± 0.008	303.2 ± 0.3	2006 May	16.15 ± 0.03
GJ 778	13.431 ± 0.011	358.1 ± 0.3	2006 May	15.34 ± 0.03
GJ 778	12.145 ± 0.007	33.0 ± 0.3	2006 May	16.37 ± 0.03
GJ 778	14.466 ± 0.007	112.3 ± 0.3	2006 May	18.18 ± 0.10
GJ 778	16.107 ± 0.005	140.6 ± 0.3	2006 May	17.78 ± 0.11
GJ 778	11.575 ± 0.008	138.8 ± 0.3	2006 May	19.48 ± 0.21
GJ 778	10.183 ± 0.011	142.3 ± 0.3	2006 May	17.93 ± 0.08
GJ 778	6.633 ± 0.012	142.4 ± 0.3	2006 May	18.42 ± 0.13
GJ 778	10.840 ± 0.011	172.1 ± 0.3	2006 May	16.81 ± 0.10
GJ 778	10.369 ± 0.013	275.4 ± 0.3	2006 May	18.36 ± 0.15
GJ 778	10.001 ± 0.010	351.6 ± 0.3	2006 May	17.32 ± 0.13
GJ 778	10.691 ± 0.010	331.1 ± 0.3	2006 May	18.97 ± 0.22
GJ 778	11.974 ± 0.007	324.2 ± 0.3	2006 May	19.44 ± 0.15

Note. — All detections represent ≥ 5 -sigma S/N levels.

^aSeparation from central star.

^bPosition angle, measured counter-clockwise from central star’s north-south axis.

*The field object described in this row is a published brown dwarf companion (Potter et al. 2002). We refrained from measuring the magnitude of this object as blending made photometry difficult.

Table 3. Target Sensitivities

Target Name	Faintest Detectable Apparent K_s -magnitude by Separation			
	1''0	2'' 0	3''0	5''0
GJ 1085	11.4	15.0	16.0	17.7
GJ 1095	10.1	13.8	15.4	17.2
GJ 159	10.7	13.5	15.1	16.9
GJ 227	11.3	14.7	15.7	17.2
GJ 230	11.9	15.3	16.6	17.8
GJ 295	12.2	15.7	16.6	17.8
GJ 311	11.7	15.3	16.3	17.8
GJ 334.2	11.3	14.7	16.0	17.4
GJ 376	11.9	15.3	16.3	17.8
GJ 407	11.6	15.0	16.3	18.0
GJ 484	11.6	15.0	16.0	17.5
GJ 547	11.2	14.7	15.7	16.3
GJ 564	12.1	15.3	16.6	17.8
GJ 56.5	11.6	15.0	16.0	17.2
GJ 611	14.0	17.5	18.2	18.7
GJ 614	11.9	15.3	16.6	17.5
GJ 651	12.5	15.1	16.1	17.8
GJ 758	8.0	11.4	12.7	14.0
GJ 778	12.5	15.7	16.9	17.8
GJ 788	11.9	15.3	16.3	17.8
GJ 895.4	11.6	15.7	16.6	17.8